







Journal of Poultry Sciences and Avian Diseases

Journal homepage: www.jpsad.com



Different light programs affect the titer of vaccination of Newcastle in broiler chicken



Alireza Dibaei¹ , Amirhossein Sharifi Moghadam² , Hadi Haghbin Nazarpak^{3*} , Nariman Sheikhi⁴ , Mahdi Askari Badouei⁵ 

¹ Faculty of Veterinary Medicine, Islamic Azad University, Garmsar Branch, Garmsar, Iran

² Faculty of Veterinary Medicine, Tehran University, Tehran, Iran

³ Department of Clinical Sciences, Faculty of Veterinary Medicine, Garmsar Branch, Islamic Azad University, Garmsar, Iran

⁴ Department of Clinical Sciences, Faculty of Veterinary Medicine, Science and Research Branch, Islamic Azad University, Tehran, Iran

⁵ Department of Pathobiology, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran

* Corresponding author email address: h.haghbin@iau-garmsar.ac.ir

Article Info

Article type:

Short Communication

How to cite this article:

Dibaei, A., Sharifi Moghadam, A., Haghbin Nazarpak, H., Sheikhi, N., & Askari Badouei, M. (2023). Different light programs affect the titer of vaccination of Newcastle in broiler chicken. *Journal of Poultry Sciences and Avian Diseases*, 1(4), 25-31.

<http://dx.doi.org/10.61186/JPSAD.1.4.25>



© 2023 the authors. Published by SANA AVIAN HOSPITAL, Tehran, Iran. This is an open access article under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License.

ABSTRACT

Light is an important factor in the functioning of the body of birds, especially the immune system. Management of the lighting program in closed systems of poultry production, mainly by affecting the concentration of melatonin, can be operational on the effectiveness of vaccination. Newcastle disease can cause extensive damage to the poultry industry. Therefore, its timely diagnosis and assessment of the success of vaccination with methods such as the hemagglutination inhibition test are of particular importance. This study investigated the effect of two photoperiods on antibody titer against Newcastle virus following vaccination. Four groups of Ross broilers were examined for six weeks. Groups A and B received the live Newcastle vaccine twice, and groups C and D were not vaccinated against Newcastle. Groups A and C received a lighting program from the management handbook of Ross Broiler, and groups B and D received the continuous lighting program (23L:1D). During the breeding period, blood samples were taken six times from each group, and the HI test measured the antibody titer. The results showed that despite the higher antibody titer in group B compared to group A, this difference was not statistically significant ($P > 0.05$). The feed conversion rate was higher in the groups that received the continuous photoperiod, and their mortality rate was lower than that of the other two groups. Comparing the results of the present study with those of various studies and articles shows the importance of the lighting regime in the success of vaccination.

Keywords: Broiler, Photoperiod, Newcastle disease, Lighting program.

Article history:

Received 20 August 2023

Revised 04 October 2023

Accepted 08 October 2023

Published online 01 December 2023

1 Introduction

Light is an important environmental factor in controlling many biological processes (1). Compared to humans, a more comprehensive range of light waves can be detected by the unique visual system of avian species (2). How light is received and processed and how it affects the biology of the bird's body is explained in detail by Davide M. Dominoni (3).

The lighting program or regime comprises three main components: wavelength, light intensity, and photoperiod. The wavelength determines the color of the light, and the photoperiod expresses the length and distribution of light and dark hours during the day and night (1, 4). In general, photoperiod is the most prominent factor in light regime management in industrial poultry farming (5). Currently, a great variety of light programs and tools are available to poultry producers, which allows them to modify light management in closed houses based on the physiological requirements of birds and breeding objectives (1, 6). For an extended period, broilers were raised under continuous photoperiods to maximize feed consumption until research showed that this light schedule caused sleep deprivation and many other problems.

In contrast, different light and dark cycles evoke the circadian rhythm of birds (5, 7). preferable functioning of the immune system, well-being of birds and reduction of aggressive behaviors, coping with stress, proper development and growth of the reproductive, skeletal, and muscular system, and obtaining more favorable production indicators are the advantages that can be achieved with the correct management of the lighting regime in industrial poultry production (1, 6, 8-10). The lighting program is important not only after hatching but also during incubation. Chickens that receive suitable light and darkness periods during the embryonic stage have better biological factors and performance during life and production (11-13).

The importance of light and the lighting program in birds' immune systems and the response to stresses has been demonstrated (4, 5). This effect is mainly attributed to the pineal gland's function in melatonin secretion (14). This hormone, primarily secreted in the dark, regulates the circadian rhythm and is generally considered an immune stimulant (15, 16). Also, in conditions of light stress, immunity can be suppressed (17). It is known that the lymphatic tissues and immune cells of birds have melatonin receptors, and their function is affected by the concentration of this hormone. The importance of this concentration in the

development of general immunity and the immunity of the respiratory system of birds (which has an undeniable role in dealing with essential diseases such as Newcastle disease and Avian Influenza) has been proven (18).

The seasonal immune change pattern can prove melatonin's immunoregulatory role in birds. In colder seasons, short day length and long darkness cause the concentration of melatonin to increase. As a result, cell proliferation and weighting in lymphatic organs such as the spleen occur. Also, in such a situation, the total number of leukocytes increases. In contrast, during warmer seasons with longer daylight hours, decreased melatonin and increased gonadal/adrenal steroid levels are responsible for decreased immune competence. Such a pattern protects the bird from the environmental stress of cold seasons and their consequences and prepares it to breed in warmer seasons (18, 19). Adequate light regimes can stimulate the expression of immune receptors (such as Toll-like receptors) (20). This positive effect on synthesizing interleukins and interferons, which are immune mediators, has also been proven (7). Appropriate photoperiod increases the mitogenic and proliferative responses of T and B lymphocytes, serum IgG level, and the percentage of CD3+, CD4+, and CD8+ cells in broilers. It has been shown that melatonin can effectively restore the impaired activity of T-helper cells and increase the activity of natural killer cells (7, 21, 22). Based on these findings, various researchers have suggested increasing the length of the dark period as an effective solution to strengthen the poultry's immune system (18, 23).

The effect of light pollution and the change of circadian rhythm on the state of the immune system of wild birds has also been investigated, which indicates a decrease in the level of immunity in these birds due to the increase in the duration of lighting (3, 24).

The results of research on the effect of the light regime and its mechanism on the immune system have been presented in more detail by other researchers (3).

Newcastle disease (ND) is caused by Avian paramyxovirus 1 (APMV-1) in many birds (including commercial poultry) (25). APMV1 infection, based on the virulence of the virus pathotype, causes a wide range of clinical manifestations, from asymptomatic viral replication to severe neurological, respiratory, and gastrointestinal symptoms and 100% mortality through sudden death (26). Due to the potential of this disease to cause significant economic losses to the poultry industry, its occurrence is reportable to the Office International des Epizooties (OIE) (27). Vaccination and biosecurity principles are

recommended as the most effective disease control strategy. Although virus isolation is considered the standard diagnostic strategy for ND, serological tests such as hemagglutination inhibition (HI) and Enzyme-Linked Immunosorbent Assay (ELISA) are the industry's first choice for diagnosing this disease (in addition to an assessment of vaccination success) (25, 28). The basis of these tests is binding the known antigen to the antibody in the serum, which evaluates the humoral immune response to the virus. Depending on the virulence of the virus, the environmental condition, and internal factors of the body, the patterns of immune response to the pathogen change (28, 29).

The present study aims to investigate the effect of a lighting program with different hours of darkness compared to a continuous lighting program in producing antibody titers against the Newcastle virus following vaccination.

2 Methods and Materials

2.1 Experimental design and husbandry conditions

Four hundred one-day-old Ross308 broiler chickens were obtained from a mother flock at 48 weeks. The levels of maternal antibodies against Newcastle disease, Avian influenza, Infectious bursal disease, and Infectious bronchitis were assessed as desirable, and the chickens were free of infection with *Mycoplasma* and *Salmonella* species. The chickens were randomly divided into four groups of 100 chickens. Each group was kept in a house that consisted of 4 parts of 2 x 5 meters with ad libitum access to feed and water for 42 days and a slaughter weight of 2-3 kg. The feed formulation was balanced according to the requirements of the Ross308 broiler and was provided to the birds in the form of pellets in three formulas: Starter (0 to 12 days old), Grower (13 to 25 days old), and Finisher (26 to 42 days old). Other breeding conditions were managed similarly for all groups based on the Ross broiler management handbook (30), and only the lighting program and receipt and non-receipt of Newcastle vaccine were different among the groups.

All four groups, at the age of 1 day and 24 days, received the H120 live vaccine against infectious bronchitis agents by eye drops and drinking water, respectively. Also, the D78 Live vaccine was administered to each group in drinking water at 15 and 22 days to protect the chickens against Infectious bursal disease. Groups A and B received the Newcastle Hitchner. B1 eye drop vaccine at the age of 8 days and the Newcastle COLON30 vaccine in drinking water on

the 18th day of rearing. Groups C and D were not vaccinated for Newcastle disease as control groups.

For groups A and C, the Ross308 breeding management light program (Table 1) was applied, and for groups B and D, a continuous light program (23L:1D) was applied.

2.1.1 Sampling and information record

On six occasions, on days 1, 7, 16, 24, 31, and 42, ten chickens were randomly selected from each group, and blood samples were taken. This procedure was performed with a 2-cc syringe without any anticoagulant agent. The collected sample was placed at room temperature for 1 hour until the serum was initially separated from the clot, and, in the next step, the serum was wholly obtained by centrifugation.

Information on the amount of feed consumed was recorded weekly, and 10% of the chickens in each group were randomly selected and carefully weighed every week. Each group's mortality and feed conversion rate (FCR) were recorded until the end of the breeding period.

2.2 Serological assessment

The sera were kept at 56°C for 30 minutes to remove the effect of non-specific inhibitors. First, 25 microliters of PBS solution were poured into all plate wells. Then, 25 microliters of the test serum were loaded in the first well, which was repeated up to the 12th well. In the next step, 25 microliters of Newcastle antigen solution with 8 HA units were added to each well. After 30 minutes, 25 microliters of red blood cell suspension of non-vaccinated chickens without maternal antibodies were added to all wells. The results were observed and recorded after 30 minutes.

2.3 Statistical analysis

Statistics were analyzed using SPSS software version 26. A chi-square test was also conducted to determine significance.

3 Results

On each occasion of serological examination, the mean, minimum, and maximum antibody titer of Newcastle disease virus in group B was higher than in group A, although statistically, this difference was not significant ($P > 0.05$) (Table 1). The maternal Newcastle antibody titer was reduced without infection in groups C and D.

On days 14, 21, 28, 35, and 42, an increasing trend was observed in the feed consumption of groups A and C. The results of weighing the chickens did not show any significant difference between the groups, but on days 14, 21, 28, and 35, a trend of weight gain was observed in groups B and D, which continued until the 42nd day.

The comparison of feed conversion rates showed that at the end of the experiment, the FCR of groups A and C was higher than groups B and D (1.85, 1.77, 1.81, and 1.75 for groups A to D, respectively). Also, the mortality rates of groups B and D were less than those of groups A and C.

Table 1. The titer of ND using HI test in Groups A and B. No significant differences between groups ($p \leq 0.05$)

Age (Day)	Group A HI Titers			Group B HI Titers		
	Min.	Mean	Max.	Min.	Mean	Max.
1	7	7	8	7	7	8
7	5	6	7	5	6.3	7
16	2	3.9	5	3	4	5
24	2	4.9	7	3	5	7
31	2	3.4	6	2	4.4	8
37	2	3.6	5	2	3.7	6
42	2	3.1	4	2	3.2	5

4 Discussion

Changes in different components of the light program in broiler breeding affect the performance of different systems of the bird's body and production results. The findings of our study revealed that rearing Ross 308 broiler chickens in a light program with more hours of light deprivation did not create a significant difference in the production of humoral immune antibodies against the Newcastle disease virus. However, the titer was numerically higher in this group.

In research conducted at the University of North Carolina in 2000, the effect of lighting conditions as an immune modulator and melatonin supplementation on the humoral and cellular immune response in Japanese quail was investigated. In the first experiment, three light treatments of short days (8:16LD), long days (16:8LD), and constant light (LL) were considered, and in the second experiment, the LL group received melatonin at different doses. Cutaneous basophil hypersensitivity reaction to phytohemagglutinin (PHA-P) as an indicator of cellular immune reaction, and primary antibody titers were evaluated seven days post-intravenous injection of Chukar red blood cell suspension were measured as an indicator of humoral immune reaction. Both 8:16LD and 16:8LD treatments had significantly better cellular and humoral immune responses than the LL group. In the second experiment, melatonin could improve the immune function of the LL group (31).

Onbaşılars *et al.* (2007) investigated the effect of two different photoperiods on production performance, occurrence of skeletal system disorders, some blood factors, and antibody production of 200 one-day-old male broiler

chicks (Ross PM3). One group received continuous lighting (24L:0D) for 42 days, and the other group experienced intermittent lighting (IL) photoperiod (1L:3D) during the same period. The results indicated that the intermittent lighting group had a higher antibody titer against NDV (32).

The study conducted by Moraes *et al.* in 2008 expressed the effect of 4 light programs of 23L:1E (23L), increasing (INC), 16L:8E (16L), and natural light (NAT) on the production and immunological performance of male Cobb-500 broiler chickens. The experiment was conducted with four groups, including 35 birds, and six repetitions. No significant difference was observed between the groups regarding antibody titer against Newcastle disease virus (33).

Gharib *et al.* 2008 evaluated the role of photoperiod and melatonin in reducing the negative impact of heat stress on broilers. In the first experiment, four groups of male Cobb × Cobb broiler chickens were raised under a continuous (23h Light: 1h Dark) and intermittent (1hL: 3hD) light program, and then one group from each light program, at a temperature of 35°C and one group was placed at a temperature of 24°C. The group that received the intermittent light program was less affected by heat stress and its adverse effects. In the second experiment, two groups were kept at 35°C, two at 24°C, and one receiving melatonin at each temperature. The group receiving melatonin showed more excellent heat stress resistance (34).

In the study conducted by Xie *et al.* in 2008, the effect of monochromatic light on the immune response of birds was investigated. Two hundred sixty-one-day-old Arbor Acres male broilers in four groups were exposed to monochromatic

red, green, blue, or white light for seven weeks. At 28 days, the group that received the green light showed a higher serum level of antibodies against the Newcastle disease virus than the group that received the red light. The level of this antibody on day 49 was higher in the group that received blue light than in the red light group. The proliferation of peripheral blood T lymphocytes increased in 21 days in green light compared to blue and red light and in 42 days in blue light compared to red light (35).

2008 Abbas *et al.* divided 300 one-day-old male broiler chicks (cobb × cobb) into three random groups. They exposed them to 3 photoperiods of continuous light (23L:1D), non-intermittent restricted light (12L:12D), and intermittent light (2L:2D) for six weeks. In chickens exposed to intermittent light, peripheral B and T lymphocyte proliferation and antibody production were significantly induced. Also, factors indicating stress, including plasma corticosterone concentration and heterophil/lymphocyte ratio, were significantly higher in the treatment 12L:12D than in the other two groups. Compared to the control group, total white blood cells (WBC) and plasma T concentration increased significantly in the intermittent light group. The results of this experiment show that intermittent photoperiods, compared to continuous or non-intermittent restricted photoperiods, improve the cellular and humoral immune function of birds and reduce stress (7).

In a study conducted by Abbas *et al.* in 2014, 150 broilers were divided into five treatments, and each treatment received a different light intensity over 42 days. This study showed that the antibody titer against the infectious bursal disease virus at six weeks of age showed a significant increase in the birds reared with the lowest light intensity compared to other groups. However, on the contrary, the antibody titer against the NDV was not significantly different between these groups at this time. Also, the Bursa of Fabricius in this group had more weight than other groups in 2 to 6 weeks (16).

In the study by Firozi *et al.* (2014), 4000 chickens were randomly placed in 4 halls and kept for 42 days with green, sunny yellow, blue, and red light. The results showed that the reduction of maternal antibodies was the lowest in the green light group. At the end of the breeding period, the birds exposed to green and blue light had the highest ND antibody titer among all groups, but this difference was not statistically significant. Birds in the yellow-light house significantly increased total serum protein compared to other groups (36).

In a study by Sharideh and Zaghari 2017, three hundred and sixty 308-day-old male chicks were reared for 42 days. Chickens were randomly divided into three groups, for which neutral white light, warm white light, and incandescent light were considered. The test results showed that the group that received warm white light had significantly higher indices of humoral immunity and antibody titers against the Newcastle disease virus than the other two groups. Although this difference was also present in the antibody titer against the Avian influenza virus, it was not statistically significant (8).

In a study conducted by Assam and Hassan in 2019, 9 different photoperiods (including 23:1, 18:6 continuous, and 18:6 intermittent photoperiods, each with three colors: red, blue, and white) for 252 Ross broiler chicken, were applied in 9 random groups during 40 days. This study showed that the 23:1 continuous photoperiod group with each light color significantly produced more antibody titers against the Newcastle disease virus than other photoperiods with the same light color (37).

In a study conducted by Tarek Mahmoud Mousa-Balabel and Karima Mohamed Abdo Abofarag in 2022, three groups of 60 commercial Indian River (IR) broiler chicks, two days before and two days after vaccination with NDV- Lasota, respectively, were treated with white light color (WLC), blue light color (BLC) and white light color and supplemented with vita E and Se in the drinking water (WES). The results showed that the body weight gain was significantly higher in broilers raised under BLC and WEC than in the WLC group. Also, these two groups had higher feed conversion ratio (FCR) and Newcastle disease virus antibody titer and lower heterophil/lymphocyte ratio compared to WLC treatment (38).

Regarding photoperiod, most of the results of other investigations were consistent with the present study's findings. However, a significant difference was observed in the antibody titer against the Newcastle virus in the experiments, which made a difference in light color. Comparing the results of various studies and articles shows the importance of the optical program and attention to all three components in the success of vaccination.

Conflict of Interest

The authors declared no conflicts of interest.

Author Contributions

H. HN. supervised the project. All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by A. D., A. S.D, M.S.B and N.S. The first draft of the manuscript was written by A. SM and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data Availability Statement

Data are available from the first author upon reasonable request.

References

1. Olanrewaju H, Thaxton J, Dozier W, Purswell J, Roush W, Branton S. A review of lighting programs for broiler production. *International journal of poultry science*. 2006;5(4):301-7. [DOI]
2. Zhang Z, Cao J, Wang Z, Dong Y, Chen Y. Effect of a combination of green and blue monochromatic light on broiler immune response. *Journal of Photochemistry and Photobiology B: Biology*. 2014;138:118-23. [PMID: 24927232] [DOI]
3. Dominoni DM. The effects of light pollution on biological rhythms of birds: an integrated, mechanistic perspective. *Journal of Ornithology*. 2015;156(Suppl 1):409-18. [DOI]
4. Zamanizad M, Ghalamkari G, Toghyani M, Adeljoo AH, Toghyani M. Effect of sequential and intermittent white, green and blue monochromatic lights on productive traits, some immune and stress responses of broiler chickens. *Livestock Science*. 2019;227:153-9. [DOI]
5. Hofmann T, Schmucker SS, Bessei W, Grashorn M, Stefanski V. Impact of housing environment on the immune system in chickens: A review. *Animals*. 2020;10(7):1138. [PMID: 32635616] [PMCID: PMC7401558] [DOI]
6. Pal P, Dey D, Sharma B, Choudhary S, Sahu J, Kumar S, Ghosh S. Effect of light management in broiler production: A review. *Journal of Entomology and Zoology Studies*. 2019;7(3):437-41.
7. Abbas AO, El-Dein AA, Desoky A, Galal MA. The effects of photoperiod programs on broiler chicken performance and immune response. *International Journal of Poultry Science*. 2008;7(7):665-71. [DOI]
8. Sharideh H, Zaghari M. Effect of light emitting diodes with different color temperatures on immune responses and growth performance of male broiler. *Annals of Animal Science*. 2017;17(2):545-53. [DOI]
9. Ryu S-T, Park B-S, Bang H-T, Kang H-K, Hwangbo J. Effects of anti-heat diet and inverse lighting on growth performance, immune organ, microorganism and short chain fatty acids of broiler chickens under heat stress. *Journal of Environmental Biology*. 2016;37(2):185.
10. Rogers AG, Pritchett EM, Alphin RL, Brannick EM, Benson ER. II. Evaluation of the impact of alternative light technology on male broiler chicken stress. *Poultry Science*. 2015;94(3):331-7. [DOI]
11. Van der Pol CW, van Roovert-Reijrink IA, Gussekloo SW, Kranenbarg S, Leon-Kloosterziel KM, van Eijk-Priester MH, et al. Effects of lighting schedule during incubation of broiler chicken embryos on leg bone development at hatch and related physiological characteristics. *Plos one*. 2019;14(8):e0221083. [PMID: 31415653] [PMCID: PMC6695123] [DOI]
12. Li X, Rathgeber B, McLean N, MacIsaac J. Providing colored photoperiodic light stimulation during incubation: 2. Effects on early posthatch growth, immune response, and production performance in broiler chickens. *Poultry Science*. 2021;100(9):101328. [PMID: 34274570] [PMCID: PMC8318990] [DOI]
13. Archer GS, Mench JA. The effects of light stimulation during incubation on indicators of stress susceptibility in broilers. *Poultry Science*. 2013;92(12):3103-8. [PMID: 24235217] [DOI]
14. Markowska M, Majewski PM, Skwarło-Sońta K. Avian biological clock-Immune system relationship. *Developmental & Comparative Immunology*. 2017;66:130-8. [PMID: 27235884] [DOI]
15. Kankova Z, Drozdova A, Hodova V, Zeman M. Effect of blue and red monochromatic light during incubation on the early post-embryonic development of immune responses in broiler chicken. *British Poultry Science*. 2022;63(4):541-7. [PMID: 35152798] [DOI]
16. Abbas G, Mahmood S, Fawwad Ahmad MY, Qureshi RA, Qamar SH, ur Rehman MZ, et al. Effect of Varying Light Intensity on Immunity Level, Dressed Weight, Minor Body Parts Weight, Fat Deposition and Serum Glucose Level in Broilers. *J Anim Vet Adv*. 2014;1:38-42.
17. Guo b Y, Li W, Chen J. Influence of nutrient density and lighting regime in broiler chickens: effect on antioxidant status and immune function. *British poultry science*. 2010;51(2):222-8. [PMID: 20461583] [DOI]
18. Verma V, Yadav S, Haldar C. Influence of environmental factors on avian immunity: an overview. *J Immun Res*. 2017;4:1028-33.
19. Haldar C, Singh S. Seasonal changes in melatonin and immunological adaptations in birds. *J Endocrinol Reprod*. 2001;5(1):13-24.
20. El-Emam H, Abou-Ismaïl U, El-Araby I, Fouda M, Ateya A. Effect of light colour temperature on expression and serum profile of selected immune markers in layers. *Journal of the Hellenic Veterinary Medical Society*. 2021;72(2):2879-88.

Acknowledgments

The authors thanks to Research deputy of the university of Garmsar Branch of Islamic Azad University, for the financial support of this research.

Ethical Consideration

The faculty of veterinary medicine, Garmsar Branch, Islamic Azad University, approved the study (code: GBIAU.VET.2020.06).

Funding Statement

This research received grant and financial support from Garmsar Branch of Islamic Azad University.

21. Kliger C, Gehad A, Hulet R, Roush W, Lillehoj H, Mashaly M. Effects of photoperiod and melatonin on lymphocyte activities in male broiler chickens. *Poultry Science*. 2000;79(1):18-25. [PMID: 10685884] [DOI]
22. Pioli C, Caroleo MC, Doriac G. Melatonin increases antigen presentation and amplifies specific and non specific signals for T-cell proliferation. *International journal of immunopharmacology*. 1993;15(4):463-8. [PMID: 8365822] [DOI]
23. Li W-b, Guo Y-l, Chen J-l, Wang R, He Y, Su D-g. Influence of lighting schedule and nutrient density in broiler chickens: effect on growth performance, carcass traits and meat quality. *Asian-Australasian Journal of Animal Sciences*. 2010;23(11):1510-8. [DOI]
24. Mishra I, Knerr RM, Stewart AA, Payette WI, Richter MM, Ashley NT. Light at night disrupts diel patterns of cytokine gene expression and endocrine profiles in zebra finch (*Taeniopygia guttata*). *Scientific reports*. 2019;9(1):15833. [PMID: 31676761] [PMCID: PMC6825233] [DOI]
25. Getabalew M, Alemneh T, Akeberegn D, Getahun D, Zewdie D. epidemiology, Diagnosis & Prevention of Newcastle disease in poultry. *Am J Biomed Sci Res*. 2019;16:50-9. [DOI]
26. Swayne DE. *Diseases of Poultry*. 14 ed: Wiley Blackwell; 2020.
27. Khabiri A, Toroghi R, Mohammadabadi M, Tabatabaeizadeh S-E, editors. Introduction of a Newcastle disease virus challenge strain (sub-genotype VII. 1.1) isolated in Iran. *Veterinary Research Forum*; 2023: Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.
28. Mao Q, Ma S, Schrickel PL, Zhao P, Wang J, Zhang Y, et al. Review detection of Newcastle disease virus. *Frontiers in Veterinary Science*. 2022;9:936251. [PMID: 35982920] [PMCID: PMC9378970] [DOI]
29. Kapczynski DR, Afonso CL, Miller PJ. Immune responses of poultry to Newcastle disease virus. *Developmental & Comparative Immunology*. 2013;41(3):447-53. [PMID: 23623955] [DOI]
30. Aviagen R. *Ross broiler management handbook*. Aviagen Limited Newbridge Midlothian EH28 8SZ, Scotland, UK. 2014.
31. Moore C, Siopes T. Effects of lighting conditions and melatonin supplementation on the cellular and humoral immune responses in Japanese quail *Coturnix coturnix japonica*. *General and Comparative Endocrinology*. 2000;119(1):95-104. [PMID: 10882554] [DOI]
32. Onbaşlılar EE, Erol H, Cantekin Z, Kaya Ü. Influence of intermittent lighting on broiler performance, incidence of tibial dyschondroplasia, tonic immobility, some blood parameters and antibody production. *Asian-Australasian Journal of Animal Sciences*. 2007. [DOI]
33. Moraes D, Lara L, Baiao N, Cancado S, Gonzalez M, Aguilar C, Lana A. Effect of lighting programs on performance, carcass yield, and immunological response of broiler chickens. *Arquivo brasileiro de medicina veterinaria e Zootecnia*. 2008;60:201-7. [DOI]
34. Gharib H, Desoky A, El-Menawey M, Abbas AO, Hendricks G, Mashaly M. The role of photoperiod and melatonin on alleviation of the negative impact of heat stress on broilers. *International Journal of Poultry Science*. 2008;7(8):749-56. [DOI]
35. Xie D, Wang Z, Dong Y, Cao J, Wang J, Chen J, Chen Y. Effects of monochromatic light on immune response of broilers. *Poultry Science*. 2008;87(8):1535-9. [PMID: 18648045] [DOI]
36. Firouzi S, Nazarpak HH, Habibi H, Jalali SS, Nabizadeh Y, Rezaee F, et al. Effects of color lights on performance, immune response and hematological indices of broilers. *J World's Poult Res*. 2014;4(2):52-5.
37. Soliman ES, Hassan RA. Impact of lighting color and duration on productive performance and Newcastle disease vaccination efficiency in broiler chickens. *Veterinary World*. 2019;12(7):1052. [PMID: 31528032] [PMCID: PMC6702559] [DOI]
38. Mousa-Balabel TM, Abofarag KMA. Blue Light Color Reduces the Newcastle Disease Post-Vaccinal Reactions of Indian River Broilers under Egyptian Conditions. 2023. [DOI]